

# Blood Transfusion Rates in Non-syndromic Craniosynostosis: A Proportion Meta-analysis of Open Versus Minimally Invasive Techniques

Mimasari Amirsyah<sup>1</sup>, Muhammad Ikhlās Abdian Putra<sup>1</sup>, Shakira Amirah<sup>2</sup>, Seba Talat Al-Gunaid<sup>3</sup>, Khalillullah Khalillullah<sup>3</sup>, Artha Maressa Theodora Simanjuntak<sup>4</sup>, Puan Faiza Nurianada<sup>5</sup>, I Nyoman Windiana<sup>6</sup>, Fawwaz Rahman Hakim<sup>7</sup>, Andi Muhammad Dzaky Tenrisessu<sup>8</sup>, Farhan Nurdiansyah<sup>7</sup> and Roy Novri Ramadhan<sup>7\*</sup>

<sup>1</sup>Plastic Reconstructive and Aesthetic Surgeon, Plastic Surgery Subdivision, Department of Surgery, Zainoel Abidin General Hospital, Banda Aceh, Indonesia

<sup>2</sup>Faculty of Medicine, Universitas Indonesia, Jakarta, Indonesia

<sup>3</sup>Faculty of Medicine, Universitas Syiah Kuala, Banda Aceh, Indonesia

<sup>4</sup>Faculty of Medicine, Universitas Gadjah Mada, Yogyakarta, Indonesia

<sup>5</sup>Faculty of Medicine, Universitas Jenderal Soedirman, Banyumas, Indonesia

<sup>6</sup>Faculty of Medicine, Universitas Pendidikan Ganesha, Buleleng, Indonesia

<sup>7</sup>Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

<sup>8</sup>Faculty of Medicine, Universitas Muslim Indonesia, Makassar, Indonesia

## ARTICLE INFO

### Article history:

Received: 6 November 2024

Accepted: 26 March 2025

### Online:

DOI 10.5001/omj.2025.67

### Keywords:

Craniosynostosis; Cranial Deformity; Blood Transfusion; Cranial Vault Remodeling; Endoscopic Surgery; Meta-analysis.

## ABSTRACT

**Objectives:** Premature suture fusion in non-syndromic craniosynostosis often requires surgery to permit normal brain growth. To minimize blood transfusion requirements, minimally invasive surgery (MIS) has been proposed. This study will pool the prevalence of blood transfusion in non-syndromic craniosynostosis surgeries, comparing open surgery (OS) with MISs. **Methods:** A systematic review and proportion meta-analysis was conducted in accordance with Preferred Reporting Items for Systematic Reviews and Meta-Analysis guidelines and registered in PROSPERO (CRD42024604632). PubMed, Scilit, Scopus, and Web of Science were searched up to 7 March 2024. Pooled prevalence was calculated using the Freeman-Tukey double arcsine transformation. **Results:** Nine studies with a total of 619 patients met the inclusion criteria. The pooled prevalence of blood transfusion was 52.0% (95% CI: 32.0–72.0) for OS and 10.0% (95% CI: 1.0–20.0) for MIS. Within OS, cranial vault remodeling exhibited the highest transfusion prevalence of 58.1% (95% CI: 37.5–78.7). Among those undergoing MIS, strip-assisted surgery had a lower prevalence (3.9%; 95% CI: -0.5–8.2) compared to endoscopic strip craniectomy (13.1%; 95% CI: -0.4–26.6). OS for metopic craniosynostosis yielded a blood transfusion prevalence of 76.2% (95% CI: 62.9–89.5), the highest among affected sutures. Sagittal craniosynostosis repairs showed a prevalence of 22.5% (95% CI: 9.6–35.3) in OS, compared with 2.6% (95% CI: 1.7–11.9) in MIS. **Conclusions:** The prevalence of blood transfusion in craniosynostosis repair is significantly lower in MIS compared to OS, suggesting the relative safety of MIS in terms of transfusion risks.

Craniosynostosis is a congenital cranial deformity resulting from the premature fusion of cranial sutures, producing abnormal head shape and risking complications such as increased intracranial pressure and disturbance on respiratory and neurologic systems.<sup>1</sup> The condition is usually non-syndromic, occurring independently from other congenital anomalies.<sup>1,2</sup> Common types include sagittal synostosis, coronal synostosis, metopic synostosis, and lambdoid synostosis.<sup>3</sup>

The primary treatment for craniosynostosis is surgical intervention, aimed at correcting the shape of

the skull and allowing for normal brain growth.<sup>4</sup> Open surgery (OS) involves a large incision that provides direct access and clear visualization of the target organs and tissues.<sup>5</sup> Principal OS techniques include cranial vault remodeling (CVR), cranial distraction osteogenesis (CDO), fronto-orbital advancement, and strip craniectomy (SC).<sup>6</sup> However, the large incisions increase the likelihood of blood transfusion, prolonged hospital stay, and post-surgical complications.<sup>7</sup>

On the other hand, minimally invasive surgery (MIS), especially endoscopic methods, requires significantly smaller incisions.<sup>8–10</sup> Its other benefits include minimum

trauma to surrounding tissues, shorter procedure time, and earlier recovery with fewer complications. In both OS and MIS, if the blood loss exceeds 100 mL (or ~10% of the patient's blood volume), blood transfusion should be administered intraoperatively.<sup>11</sup>

Measuring the prevalence of blood transfusion is crucial for perioperative planning including blood bank logistics, patient safety, and resource allocation.<sup>11,12</sup> Existing studies are mostly single-center, with limited sample sizes.<sup>12–15</sup> To derive robust estimates of blood transfusion rates in craniosynostosis surgeries, we conducted this multicenter proportion meta-analysis.<sup>16–19</sup>

## METHODS

The study employed a proportional meta-analysis design, as per the Preferred Reporting Items for Systematic Reviews and Meta-Analysis protocol.<sup>20</sup> The research question was, “What is the prevalence of blood transfusion in OS and MOS for the management of non-syndromic craniosynostosis?” The study was officially recorded in PROSPERO (registration number CRD42024604632, November 01, 2024).

On 7 March 2024, we queried PubMed, Scilit, Scopus, and Web of Science databases using the following search terms: “craniosynostosis” OR “non-syndromic craniosynostosis” OR “cranial suture synostosis” AND “open surgery” OR “cranial vault remodeling” OR “strip craniectomy” OR “endoscopic surgery” OR “minimally invasive surgery” AND “blood transfusion” OR “transfusion requirement” OR “perioperative blood loss” OR “intraoperative transfusion”. We used the recommended search syntax and rules specific to each database.<sup>21</sup>

**Inclusion criteria:** Studies focusing on patients with non-syndromic craniosynostosis undergoing OS or MIS, including CVR, endoscopic SC (ESC), spring-assisted surgery (SAS), SC, remodeling with helmet therapy (RH), and posterior vault distraction (PiP). Eligible studies were required to report the prevalence or incidence of blood transfusion as an outcome measure. Accepted study designs included cross-sectional, case-control, and cohort studies that provided primary patient data relevant to surgical outcomes.

**Exclusion criteria** included studies with non-comparative or incomplete data, case series without distinct outcome reporting, review articles, commentaries, editorials, and conference abstracts.

Following automatic duplicate removal using EndNote 19, screening was carried out in two phases: first by evaluating titles and abstracts, and then by reviewing full texts. Two authors independently conducted each stage, resolving disagreements through consensus; if consensus could not be reached, a third author was consulted for clarification. The Newcastle-Ottawa Scale was employed for the critical appraisal of observational studies, which suggests a maximum score of 9 for cohort studies and 7 for cross-sectional studies.<sup>22,23</sup>

We extracted the following data from each study: first author, year of publication, country, study design, sample size (mean patient age in months, sex distribution), suture type (sagittal, coronal, metopic, lambdoid, or mixed), surgical technique, and blood transfusion rates. Continuous variables were reported as mean  $\pm$  SD, while other data were converted using standard formulae.<sup>24–26</sup> Data extraction was carried out by one author and verified and confirmed by another author.

Meta-analysis was performed using Rstudio version 2024.04.2 (Posit, PBC, Boston, MA, USA). The analysis focused on both OS (such as CVR and SC) and MIS (including spring-assisted and spring-endoscope surgeries). The pooled proportion was estimated by Freeman-Tukey double arcsine transformation (FTT) method. Back-transformation was performed based on the inverse of the variance of the pooled Freeman-Tukey double arcsine transformation proportion. Prevalence was calculated by multiplying 100% with the back-transformed value. Heterogeneity among the studies was evaluated by  $I^2$  statistic, with cut-off criteria of 50% and  $p$ -Het < 0.1 indicating high heterogeneity. Subgroup analysis compared surgical techniques (CVR, SC, SAS, ESC).


## RESULTS

A total of 1473 records were retrieved. After removing 904 duplicates, 569 records were screened. Full texts for 102 articles were retrievable; these were scrutinized based on the eligibility criteria. Six were duplicates, and 78 studies failed our eligibility criteria due to wrong design (8), missing outcome data (64), ineligible study designs (6), or did not exclude syndromic patients (4).<sup>27–30</sup> Finally, nine studies were identified as eligible to include in our systematic review and meta-analysis [Table 1]. The flow-diagram of the screening and selection process is presented as Figure 1.

**Table 1:** Characteristics and qualities of studies eligible for the estimation of blood transfusion prevalence in craniosynostosis repairs.

| Study                              | Location       | Study design    | Affected suture(s) | Technique | Group | Category | Total | Age, months | Sex, M/F | Blood transfusion, % | Quality | Professionals                                   |
|------------------------------------|----------------|-----------------|--------------------|-----------|-------|----------|-------|-------------|----------|----------------------|---------|-------------------------------------------------|
| Badiee et al, 2022 <sup>14</sup>   | United States  | Cross-sectional | Metopic            | SC        | OS    | OS       | 15    | 5.5 ± 1.7   | 14/1     | 53.3                 | 6       | 1 neurosurgeon                                  |
|                                    |                |                 |                    | CVR       |       |          | 20    | 9.8 ± 0.8   | 14/6     | 85.0                 |         | 4 craniofacial surgeons + 1 neurosurgeon        |
| Kamel et al, 2021 <sup>31</sup>    | United States  | Cross-sectional | Coronal            | FOA       | OS    | OS       | 64    | 11.6 ± 10.8 | 19/45    | 80.0                 | 7       | 2 neurosurgeons + 2 craniofacial surgeons       |
|                                    |                |                 |                    | CDO       |       |          | 17    | 9 ± 1.6     | 3/14     | 100                  |         | 1 craniofacial surgeon + 2 neurosurgeons        |
| Magge et al, 2019 <sup>32</sup>    | United States  | Cross-sectional | Sagittal           | CVR       | OS    | OS       | 21    | 5.1 ± 2.7   | NR       | 33.3                 | 7       | Multiple surgeons                               |
|                                    |                |                 |                    | ES        |       |          | 30    | 3.1 ± 1.2   | NR       | 16.7                 |         | Multiple surgeons                               |
| Jivraj et al, 2019 <sup>33</sup>   | United Kingdom | Cohort          | Unicoronal         | FO-CVR    | OS    | OS       | 23    | 18.5 ± 4.2  | 10/13    | 60.9                 | 8       | Not applicable                                  |
| Isaac et al, 2018 <sup>34</sup>    | United States  | Cross-sectional | Coronal            | SC        | OS    | OS       | 115   | 3.3 ± 2.0   | 50/65    | 1.7                  |         |                                                 |
|                                    |                |                 |                    | FOA       | OS    | OS       | 34    | 10.4 ± 2.3  | NR       | 30.0                 | 7       | 1 neurosurgeon + 2 plastic surgeon              |
|                                    |                |                 |                    | ES        | MIS   | MIS      | 60    | 3.0 ± 1.1   | NR       | 0.0                  |         | 1 neurosurgeon                                  |
| Melin et al, 2020 <sup>7</sup>     | United States  | Cohort          | Mix                | CVR       | OS    | OS       | 18    | < 6         | NR       | 90.0                 | 9       | 1 neurosurgeon + 1 craniofacial plastic surgeon |
|                                    |                |                 |                    | ES        | MIS   | MIS      | 33    | < 6         | NR       | 42.0                 |         | 1 neurosurgeon + 1 craniofacial plastic surgeon |
| Mendonca et al, 2020 <sup>15</sup> | India          | Cohort          | Sagittal           | ES        | MIS   | MIS      | 17    | 3.8 ± 0.6   | 4/13     | 0.0                  | 9       | Not applicable                                  |
|                                    |                |                 |                    | CVR       | OS    | OS       | 23    | 11.9 ± 2.6  | 10/13    | 27.0                 |         |                                                 |
| Smith et al, 2021 <sup>35</sup>    | United States  | Cohort          | Sagittal           | SC        | OS    | OS       | 17    | 12.6 ± 3.2  | 14/3     | 11.8                 | 9       | 5 neurosurgeons + 1 plastic surgeon             |
| Skolnic et al, 2020 <sup>36</sup>  | United States  | Cross-sectional | Sagittal           | SAS       | MIS   | MIS      | 45    | 19.9 ± 3.1  | 33/12    | 11.1                 |         | 5 neurosurgeons                                 |
|                                    |                |                 |                    | SAS       | MIS   | MIS      | 27    | 4.5 ± 1.3   | 23/4     | 0.0                  | 7       | 1 neurosurgeon                                  |

NR: not reported; SC: strip craniectomy; CVR: cranial vault remodeling; FOA: fronto-orbital advancement; CDO: cranial distraction osteogenesis; ES: endoscopic surgery; FO-CVR: fronto-orbital cranial vault remodeling; SAS: strip-assisted surgery; ES-HM: endoscopic surgery with helmet molding




**Figure 1:** Flow-diagram for the screening and selection of the studies eligible for the estimation of blood transfusion prevalence in craniosynostosis repairs.

Table 1 summarizes the nine studies. Seven were from the US, one from the UK, and one from India. Together they had a total of 619 patients aged 3–19.9 years. The quality of eight studies was assessed as ‘good’

using Newcastle–Ottawa Scale. One study was assessed ‘moderate’ due to small sample size (n = 35).<sup>14</sup> Another concern was found in Jivraj et al,<sup>31</sup> because their control group data was sourced from external medical records, thus affecting comparability.<sup>31</sup>

The pooled prevalence of blood transfusion for OS or MIS are presented in Figure 2 (a and b). The prevalence for the OS was 52.0% (95% CI: 32.0–72.0). As for the MIS, the prevalence was 10.0% (95% CI: 1.0–20.0). Heterogeneities were high in both pooled estimates with  $I^2$  values of 92.1% ( $p < 0.001$ ) and 97.5% ( $p < 0.001$ ) for OS and MIS, respectively. Egger’s correlation suggested publication bias ( $p < 0.001$ ) in the pooled analysis of blood transfusion prevalence among patients undergoing OS, but negligible bias for MIS ( $p = 0.955$ ). The symmetry and asymmetry of the funnel plot data are presented in Figure 2 (c and d).

Results from the pooled estimates based on the surgical techniques and types of sutures are presented in Table 2. The highest blood transfusion prevalence was observed in CDO (100%), though it was only reported in one study. The second highest prevalence was observed in CVR (58.1%; 95% CI: 37.5–78.7), while SC (10.5%; 95% CI: 1.4–19.5) occupied the last position among the OS. The lowest prevalence among MIS was found in SAS (3.9%; 95% CI: -0.5–8.2), followed by ES with a prevalence of 13.1% (95% CI: -0.4–26.6). Among OS, from the highest to the lowest



**Figure 2:** (a) Pooled prevalence of blood transfusion among patients undergoing open and (b) minimally invasive surgeries. (c) The funnel plot for publication bias detected in the pooled analysis of the prevalence analysis in open and (d) minimally invasive surgery groups.


**Table 2:** Subgroup analysis for the pooled estimates of blood transfusion prevalence in craniosynostosis repairs.

| Variable                     | No of Studies | Sample size, n | Prevalence, % (95% CI) | <i>I</i> <sup>2</sup> , % | <i>p</i> -Het  |
|------------------------------|---------------|----------------|------------------------|---------------------------|----------------|
| Open surgeries               |               |                |                        |                           |                |
| Cranial vault remodeling     | 7             | 203            | 58.1 (37.5–78.7)       | 92.0                      | <0.001*        |
| Strip craniectomy            | 3             | 147            | 10.5 (1.4–19.5)        | 92.1                      | <0.001*        |
| Osteogenesis                 | 1             | 17             | 100 (not applicable)   | Not applicable            | Not applicable |
| Minimally invasive surgeries |               |                |                        |                           |                |
| Spring-assisted              | 2             | 72             | 3.9 (-0.5–8.2)         | 67.6                      | 0.079          |
| Spring-endoscope             | 5             | 180            | 13.1 (-0.4–26.6)       | 93.1                      | 0.057          |
| Suture                       |               |                |                        |                           |                |
| Open surgeries               |               |                |                        |                           |                |
| Coronal                      | 3             | 115            | 69.2 (29.8–10.9)       | 97.7                      | <0.001*        |
| Sagittal                     | 3             | 61             | 22.5 (9.6–35.3)        | 37.1                      | 0.210          |
| Metopic                      | 2             | 35             | 76.2 (62.9–89.5)       | 77.1                      | 0.037*         |
| Unicoronal                   | 2             | 138            | 2.6 (0.2–4.9)          | 97.0                      | <0.001*        |
| Minimally invasive           |               |                |                        |                           |                |
| Sagittal                     | 5             | 159            | 2.6 (1.7–11.9)         | 48.2                      | 0.103          |

\*Significant

prevalence was observed in metopic (76.2%; 95% CI: 62.9–89.5), coronal (69.2%; 95% CI: 29.8–10.87), sagittal (22.5%; 95%CI: 9.6–35.3), and unicoronal sutures (2.6%; 95% CI: 0.2–4.9). In the case of MIS, we were only able to pool the data for sagittal sutures, where the prevalence was found to be 2.6% (95% CI: 1.7–11.9).

Figure 3 presents the forest plots for the pooled prevalence of blood transfusion in various craniosynostosis repair techniques. The prevalence varied according to the surgical technique. Open CVR demonstrated the highest prevalence at 58.1% (95% CI: 37.5–78.7), while SC, spring-assisted surgery, and endoscopic-assisted surgery had significantly lower rates (4–20%).

**Figure 3:** Forest plots for the pooled prevalence of blood transfusion in various craniosynostosis repair techniques such as (a) open cranial vault remodeling, (b) strip craniectomy, (c) spring-assisted surgery, and (d) endoscope-assisted surgery.

Among open techniques, CVR demonstrated the highest transfusion prevalence, with pooled rates approaching 58% (95% CI: 37–79) [Figure 3a]. SC showed a lower transfusion prevalence of 20% (95% CI: -9–49) [Figure 3b]. SAS had the lowest transfusion prevalence at only 4% (95% CI: 0–8), suggesting minimal transfusion requirements [Figure 3c]. Endoscopic-assisted surgery showed a slightly higher pooled prevalence of 13% (95% CI: 0–27), but still substantially lower than traditional open techniques [Figure 3d].

## DISCUSSION

Variables such as patient age, blood volume dynamics, and surgical complexity influence transfusion risk in craniosynostosis repair. Although infants < 24 months are reported to have lower infection rates and shorter hospital stays, their limited physiological reserve increases susceptibility to intraoperative blood loss. Consequently, the overall need for blood transfusion does not significantly differ across age groups.<sup>32–34</sup> However, younger patients may require prolonged hospitalization due to the need for closer postoperative monitoring.<sup>35</sup> Importantly, only a minority of transfusion-free hospitalizations have been documented after craniofacial reconstruction.<sup>33</sup> The timing of surgery also significantly influences blood loss and transfusion rates. Optimal intervention within the first year of life, particularly in syndromic cases, reduces complications and supports neurodevelopmental outcomes, whereas delayed intervention is associated with higher perioperative risks.<sup>36</sup> Though Blood loss is generally proportional to duration of surgery,<sup>37</sup> reported transfusion rates vary widely (20–100%).<sup>38</sup> In severe cases, rapid hemorrhage may necessitate massive transfusion.<sup>38,39</sup>

Transfusion management in craniosynostosis requires a patient-specific strategy that considers surgical technique, intraoperative hemodynamics, and transfusion thresholds. OS often necessitate early transfusion, whereas minimally invasive procedures typically require less. Guidelines recommend transfusion when hemoglobin falls < 8 g/dL in stable patients or 10 g/dL in unstable patients, with special considerations for infants with congenital conditions.<sup>40</sup> Effective management must balance evidence-based thresholds with patient physiology and intraoperative judgment to minimize complications. Open procedures usually involve extensive tissue manipulation and larger

incisions.<sup>37,39</sup> This indicates that surgical approach and suture involvement, rather than age alone, are key determinants of transfusion needs. Our meta-analysis demonstrated that MIS techniques substantially reduce transfusion requirements, supporting their prioritization in patients at higher risk of transfusion-related complications or in settings with limited blood supply.<sup>33</sup> Furthermore, transfusion prevalence data can inform perioperative planning and optimize resource allocation. Of note, OS in the included studies were performed mainly by neurosurgeons and craniofacial plastic surgeons, whereas only neurosurgeons performed MIS. Age distribution also differed, with OS typically performed in older infants (~9–17 months) and MIS in younger infants (~3–5 months).

While guidelines recommend transfusion initiation < 10 g/dL in stable pediatric patients, emerging evidence suggests that restrictive transfusion strategies may even reduce mortality in non-craniosynostosis surgeries.<sup>38</sup> Such findings underscore the importance of tailoring transfusion practices to patient-specific and procedural factors. Prioritizing procedures with lower transfusion requirements also facilitate cost and resource savings.<sup>36</sup> The choice of surgical technique, however, extends beyond transfusion risk. A prior survey indicated that selection often depends on skull maturation.<sup>32</sup> For infants older than 12 months, more invasive approaches such as CVR and CDO have been preferred, as extensive exposure allows more complete correction of cranial deformities.<sup>33</sup> Neurodevelopmental implications also warrant consideration in surgical planning.<sup>34,36</sup>

Taken together, these findings support broader implementation of MIS techniques where feasible, especially in resource-limited hospitals or in patients at high transfusion risk. Improved training opportunities and allocation of resources for MIS are recommended.

The main limitation of this study is the high heterogeneity across prevalence estimates. However, this variability highlights the need for tailored transfusion strategies in craniosynostosis repair. Caution should be applied when generalizing these findings, as institutional protocols and local resources may strongly influence transfusion practices. Large-scale, multicenter studies are needed in the future to compare long-term outcomes of OS versus MIS, not only regarding transfusion but also recurrence rates, neurodevelopment, and patient quality of life. Standardized study designs will be essential to refine surgical guidelines and optimize perioperative blood management.

## CONCLUSION

There are significant differences in blood transfusion requirements between OS and MIS for craniosynostosis repair. The pooled prevalence of blood transfusion is notably higher in OS (52.0%) than in MIS (10.0%). Procedures such as CVR and metopic suture repairs are associated with particularly high transfusion rates. Our findings suggest that MIS techniques may minimize transfusion requirements, and improve patient safety and outcome. Targeted blood management strategies are recommended for high-risk OS cases.

### Disclosure

The authors declare no conflicts of interest. No funding was received for this study.

### REFERENCES

- Blessing M, Gallagher ER. Epidemiology, genetics, and pathophysiology of craniosynostosis. *Oral Maxillofac Surg Clin North Am* 2022 Aug;34(3):341-352.
- Spazzapan P, Velnar T. Isolated sagittal craniosynostosis: a comprehensive review. *Diagnostics (Basel)* 2024 Feb;14(4):435.
- Ornoy A. Craniofacial malformations and their association with brain development: the importance of a multidisciplinary approach for treatment. *Odontology* 2020 Jan;108(1):1-15.
- Xue AS, Buchanan EP, Hollier LH Jr. Update in management of craniosynostosis. *Plast Reconstr Surg* 2022 Jun;149(6):1209e-1223e.
- Villavisanis D, Blum JD, Cho DY, Barrero C, Swanson JW, Nah H-D, et al. Suture fusion, cephalic index, and head shape in non-syndromic sagittal craniosynostosis. *Plast Reconstr Surg Glob Open* 2022;10:132-133.
- Shruthi NM, Gulati S. Craniosynostosis. *J Pediatr Neurosci* 2022 Sep;17(Suppl 1):S54-S60.
- Melin AA, Moffitt J, Hopkins DC, Shah MN, Fletcher SA, Sandberg DI, et al. Is less actually more? An evaluation of surgical outcomes between endoscopic suturectomy and open cranial vault remodeling for craniosynostosis. *J Craniofac Surg* 2020 Jun;31(4):924-926.
- Al-Shaqsi SZ, Lam NW, Forrest CR, Phillips JH. Endoscopic versus open total vault reconstruction of sagittal craniosynostosis. *J Craniofac Surg* 2021 May;32(3):915-919.
- Albuz B, Coskun ME, Egemen E. Endoscopy-assisted craniosynostosis surgery versus cranial vault remodeling for non-syndromic craniosynostosis: experience of a single center. *Türk Neurosurg* 2024;34(1):102-112.
- Porto Junior S, Meira DA, da Cunha BL, Fontes J, Pustilnik HN, da Silva da Paz MG, et al. Endoscopic surgery for craniosynostosis: A systematic review and single-arm meta analysis. *Clin Neurol Neurosurg* 2024 Jul;242:108296.
- Hevia-Rodríguez P, Romero-López C, Martín-Fernández J, Rivero-Garvía M, Márquez-Rivas J. Endoscopy-assisted craniosynostosis surgery without postoperative helmet molding therapy. *World Neurosurg* 2024 Mar;183:79-85.
- Kurlander DE, Ascha M, Marshall DC, Wang D, Ascha MS, Tripi PA, et al. Impact of multidisciplinary engagement in a quality improvement blood conservation protocol for craniosynostosis. *J Neurosurg Pediatr* 2020 Jun;26(4):406-414.
- Suarez AD, Taicher B, Fuchs H, Marcus J, Vestal M, Homi M, et al. Predictors of blood transfusion for endoscopic assisted craniosynostosis surgery. *J Craniofac Surg* 2022 Jul-Aug;33(5):1327-1330.
- Badiee RK, Popowitz E, Mark IT, Alcon A, Hwang J, Rorison E, et al. Superior long-term appearance of strip craniectomy compared with cranial vault reconstruction in metopic craniosynostosis. *Plast Reconstr Surg Glob Open* 2022 Feb;10(2):e4097.
- Mendonca DA, Gopal S, Gujjalanavar R, Deraje V, H.R. S, Ramamurthy V. Endoscopic versus open cranial reconstruction surgery for anterior craniosynostosis: experience from South-East Asia. *FACE* 2020;1:105-113.
- Putra MI, Gusti N, Duta TF, Alina M, Qanita I, Naufal MA, et al. Vitamin D supplementation improves foot ulcers among diabetic patients: Pooled analysis of randomized controlled trials. *Narra X* 2023;1(3): e104.
- Refin RY, Andika FF, Abudurrahman MF, Maidar M, Yufika A, Mulya IC, et al. Can smartphone-based diabetes control apps improve cardiovascular risk among patients with diabetes? A systematic review and meta-analysis. *Narra X* 2024;2:e123.
- Wikurendra EA, Aulia A, Fauzi ML, Fahmi I, Amri I. Willingness-to-pay for urban green space: A meta-analysis of surveys across China. *Narra X* 2023 Dec;1(3).
- Sandeep M, Padhi BK, Yella SS, Sruthi KG, Venkatesan RG, Krishna Sasanka KB, et al. Myocarditis manifestations in dengue cases: A systematic review and meta-analysis. *J Infect Public Health* 2023 Nov;16(11):1761-1768.
- Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021 Mar;372(71):n71.
- Putra MI, Amirsyah M, Budiman B, Amirah S, Al-Gunaid ST, Iqhrammullah M. Comparison of surgical techniques in managing craniosynostosis: systematic review and bayesian network meta-analysis. *Surgeries (Basel)* 2024;5:970-985.
- Iqhrammullah M, Gusti N, Andika FF, Abdullah A. Association of serum vitamin D and the risk of cardiovascular diseases among diabetic patients: A systematic review and meta-analysis. *Clin Nutr ESPEN* 2024 Aug;62:66-75.
- Iqhrammullah M, Duta TF, Alina M, Qanita I, Naufal MA, Henira N, et al. Role of lowered level of serum vitamin D on diabetic foot ulcer and its possible pathomechanism: A systematic review, meta-analysis, and meta-regression. *Diabetes Epidemiol Manag* 2024;13:100175.
- Shi J, Luo D, Wan X, Liu Y, Liu J, Bian Z, et al. Detecting the skewness of data from the five-number summary and its application in meta-analysis. *Stat Methods Med Res* 2023 Jul;32(7):1338-1360.
- Luo D, Wan X, Liu J, Tong T. Optimally estimating the sample mean from the sample size, median, mid-range, and/or mid-quartile range. *Stat Methods Med Res* 2018 Jun;27(6):1785-1805.
- Shi J, Luo D, Weng H, Zeng XT, Lin L, Chu H, et al. Optimally estimating the sample standard deviation from the five-number summary. *Res Synth Methods* 2020 Sep;11(5):641-654.
- Park H, Choi JW, Ra YS. One-piece fronto-orbital distraction osteogenesis without bandeau in patients with coronal craniosynostosis: A five-year follow-up retrospective study of 45 consecutive patients. *J Craniomaxillofac Surg* 2022 Dec;50(12):884-893.
- Massenburg BB, Nassar AH, Hopper RA. National database reported outcomes following craniosynostosis reconstruction. *J Craniofac Surg* 2020;31(1):154-157.
- Hoppe IC, Taylor JA. A cohort study of strabismus rates following correction of the unicoronal craniosynostosis deformity: Conventional bilateral fronto-orbital advancement versus fronto-orbital distraction osteogenesis. *J Craniofac Surg* 2021 Oct;32(7):2362-2365.
- Kanack MD, Proctor MR, Meara JG, Balkin DM, Rodean J, Stringfellow IC, et al. Hospital variation and resource use for infants with craniosynostosis undergoing

- open, endoscopic, and distraction osteogenesis surgical techniques. *Cleft Palate Craniofac J* 2025 May;62(5):845-852.
31. Jivraj BA, Ahmed N, Karia K, Menon R, Robertson E, Sodha A, et al. A 24-month cost and outcome analysis comparing traditional fronto-orbital advancement and remodeling with endoscopic strip craniectomy and molding helmet in the management of unicoronal craniosynostosis: A retrospective bi-institutional review. *JPRAS Open* 2019 Feb;20:35-42.
  32. Puthumana JS, Lopez CD, Lake IV, Yang R. Evaluation of complications and outcomes in craniosynostosis by age of operation: Analysis of the national surgical quality improvement program-pediatric. *J Craniofac Surg* 2023 Jan-Feb;34(1):29-33.
  33. Beethe AB, Spitznagel RA, Kugler JA, Goeller JK, Franzen MH, Hamlin RJ, et al. The road to transfusion-free craniosynostosis repair in children less than 24 months old: A quality improvement initiative. *Pediatr Qual Saf* 2020 Jul;5(4):e331.
  34. Puente-Espel J, Rios-Lara y Lopez RL, Moreno-Álvarez MC, Morel-Fuentes EJ. Craniosynostosis: A multidisciplinary approach based on medical, social and demographic factors in a developing country. *Rev Med Hosp Gen (Mex)* 2016;79:230-239.
  35. van Uitert A, Megens JH, Breugem CC, Stubenitsky BM, Han KS, de Graaff JC. Factors influencing blood loss and allogeneic blood transfusion practice in craniosynostosis surgery. *Paediatr Anaesth* 2011 Dec;21(12):1192-1197.
  36. Bruce MR, Frasco PE, Sell-Dortin KA, Cuevas CV, Chang Y-H, Lim ES, et al. Days alive and out of the hospital after heart transplantation: A retrospective cohort study. *J Cardiothorac Vasc Anesth* 2024 Jan;38(1):93-100.
  37. Ali Z, Hassan N, Syed S. Blood transfusion practices in neuroanaesthesia. *Indian J Anaesth* 2014 Sep;58(5):622-628.
  38. Kumba C. Hemoglobin levels and postoperative outcome in pediatric surgical patients. *SOJ Pediatr Clin Neonato* 2021;1(3):1-8.
  39. Park S-Y, Seo K-S, Karm M-H. Perioperative red blood cell transfusion in orofacial surgery. *J Dent Anesth Pain Med* 2017 Sep;17(3):163-181.
  40. Carson JL, Stanworth SJ, Guyatt G, Valentine S, Dennis J, Bakhtary S, et al. Red blood cell transfusion: 2023 AABB international guidelines. *JAMA* 2023 Nov;330(19):1892-1902.